

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2006		2. REPORT TYPE		3. DATES COVERED 00-00-2006 to 00-00-2006	
4. TITLE AND SUBTITLE Finite Element-Based Coastal Ocean Modeling: Today and Tomorrow				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory, Oceanography Division, 1002 Balch Boulevard, Stennis Space Center, MS, 39522-5001				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Finite Element-Based Coastal Ocean Modeling: Today and Tomorrow

C.A. Blain, T.C. Massey, R.A. Arnone, and R.W. Gould
Oceanography Division

Introduction: The continued necessity of military special forces operations in riverine and coastal environments along with increasing civilian concerns related to sediment transport, search and rescue, pollutant dispersal, and coastal restoration, have resulted in the need for detailed knowledge of currents and water levels in coastal, estuarine, and riverine environments. This demand for information at highly resolved spatial and temporal scales and the availability of massively parallel computer resources has brought to the forefront the capabilities of finite element (FE)-based coastal ocean circulation models. The use by these models of unstructured triangular meshes permits a large degree of flexibility in representing the complexities of coastal environments associated with convoluted shorelines, and steep gradients in currents or bathymetry. Ultimately this flexibility results in model predictions over periods of hours or less at spatial scales that range from meters to kilometers.

A Model for Coastal Mississippi and Louisiana: Unstructured meshes containing upwards of 281,800 computational points and 540,000 triangular elements are constructed to represent riverine and coastal currents off the shores of southeast Louisiana and Mississippi (Fig. 1). Spatial resolution is 50 m in rivers and decreases to 100 to 200 m offshore. The need for remote boundary forcing is accommodated by including the Gulf of Mexico and extending the mesh into the north Atlantic ocean. The applied forcing includes tides, wind stress at the water surface and river discharge from the Atchafalaya and Mississippi Rivers.

Connecting Mississippi River Currents and Observed Optical Properties: The underlying physical processes that influence observed optical distributions as derived from remotely-sensed ocean color data are often too complex and are manifest at space and time scales that preclude full understanding from imagery alone. But when coupled with highly resolved, dynamically advanced numerical models, understanding increases and the prediction of dynamical processes and optical patterns in coastal waters is possible. During March 2004, magnitudes of the depth-averaged currents computed at the mouth of the




FIGURE 1

Detail of the unstructured computational mesh constructed for Louisiana and Mississippi coastal waters. The mesh, containing 281,528 computational points and 455,203 triangular elements, has 50-m resolution of the Mississippi and Atchafalaya rivers, 100-m resolution of nearshore areas, and includes the entire Gulf of Mexico extending to 60 deg W in the north Atlantic ocean to accommodate remote tidal forcing from a global tide model.

Mississippi River and westward are correlated to the spatial distribution of the beam attenuation coefficient (a sum of the total absorption and scattering) derived from the MODIS 250-m ocean color data.¹ The plume of high beam attenuation (an indication of more sediment particles in the water) found westward of the Mississippi River outflow is a response to the stronger, undulating westward moving coastal currents (Fig. 2). Comparisons of the same currents with observed chlorophyll are not as striking; this is expected since many unmodeled biological and optical processes contribute to the distribution of chlorophyll in coastal waters. Remotely sensed ocean color data are often the only observational window on ocean processes in denied areas of military interest, making the coupling of satellite imagery with numerical models imperative for developing a predictive capability in such regions.

Predicting Hurricane Katrina Storm Surge:

Tidal variability combined with the best available NOAA reanalyzed surface winds representing the evolution of Hurricane Katrina force the FE-based coastal model in a hindcast of the surge elevation experienced along the Mississippi Gulf coast. Model predictions (Fig. 3, top) at landfall on August 29, 2005, 5:30Z indicate maximum surges are in the range of 9 m and

occur several hours after the peak winds (Fig. 3, lower panels). The unstructured mesh includes such features as levees along the Mississippi River, channels, inland topography, and a capability for surge overtopping of barrier islands.

A New Approach for the Future: While the current FE methodology leads to robust computations of tide and surge dynamics, limitations on local conservation properties are motivating development of the next generation of FE-based coastal models. NRL is implementing a new and promising class of FE methods, the discontinuous Galerkin (DG) methods, which have seen very rapid development over the last several years. The primary advantage of DG methods is their local (element by element) enforcement of the conservation laws, which pave the way for accurate transport of temperature and salinity, or other conservative tracers. Findings from comparisons between the existing FE-based model and a model using the new DG methodology² show that the DG formulation is able to retain the robustness of existing model solutions while more sharply resolving gradients and other small-scale circulation features. Furthermore, the local nature of the DG method is advantageous in permitting 1) elemental specification of the order




FIGURE 2

The beam attenuation coefficient derived from the MODIS 250-m ocean color data (left) is well correlated to 2D finite element model-predicted barotropic currents (right) on March 17, 2004. The westward movement of the sediment plume (high beam attenuation/high particle concentration) from the Mississippi River can be attributed in part to the underlying westward movement and undulation of the currents.




FIGURE 3
Hindcast of Hurricane Katrina storm surge elevations near the time of landfall August 29, 2005, 5:30Z. Time series evolutions of the wind and surge at Waveland, MS and Pass Christian, MS.

of approximation, thereby allowing for the capture of frontal and sharp gradient regions, 2) nonconforming unstructured meshes that simplify development of adaptive meshing strategies, and 3) a reduction in the computational footprint to an element and immediate neighbor elements, a necessity for efficient parallel processing on distributed memory machines. The potential of this new methodology to revolutionize coastal ocean modeling is significant and NRL is leading the way in its development and application to complex ocean dynamics problems.

Acknowledgments: The authors recognize Dr. Joannes Westerink at the University of Notre Dame for

contributing an early version of the developed finite element mesh and Dr. Frank Giraldo of NRL's Marine Meteorology Division for his contributions to the future DG-based model.

[Sponsored by NRL]

References

- ¹ R.A. Arnone, Z.P. Lee, P. Martinovich, and S.D. Ladner, "Characterizing the Optical Properties of Coastal Waters by Coupling 1 km and 250 m Channels on MODIS - Terra," *Ocean Optics XVI*, 2002.
- ² C.A. Blain and T. C. Massey, "Application of a Coupled Discontinuous-Continuous Galerkin Finite Element Shallow Water Model to Coastal Ocean Dynamics," *Ocean Modeling* **10**, 283-315, 2005. ★